

Registered Name:	Trucker's Freak On A Leash	Owner:	Sanna Koponen
Nickname:	Bruno	Country:	Finland
Registration ID:	ER47005/19	Testing date:	2019/10/28
Microchip:	985141001208561		Identified with standard
Breed:	Cirneco dell'Etna	identification profile:	ISAG 2006 markers
Gender:	Male	prome.	

Dog's identity verified from microchip or tattoo by veterinarian or other authorised person during sample taking: Yes

# Test results for pharmacogenetics

Disorder	Mode of Inheritance	Result
Multi-Drug Resistance 1, (MDR1)	Autosomal Dominant	Clear

On behalf of Genoscoper Laboratories,

boner ma SIGNATURE



 Registered Name:
 Trucker's Freak On A Leash
 Owner:
 Sanna Koponen

 Nickname:
 Bruno
 Country:
 Finland

 Registration ID:
 ER47005/19
 Testing date:
 2019/10/28

 Microchip:
 985141001208561
 DNA
 Identified with standard identification profile:

 Breed:
 Cirneco dell'Etna
 profile:
 JSAG 2006 markers profile:

Dog's identity verified from microchip or tattoo by veterinarian or other authorised person during sample taking: Yes

### Test results - Traits - page 1

#### Coat Type

Trait	Genotype	Description
Coat Length	L/L	The dog is likely to have short-haired coat.
Furnishings / Improper Coat in Portuguese Water Dogs (marker test)	GG/CC	The dog is not genetically likely to express furnishings.
<i>KRT71</i> c.451C>T (p.Arg151Trp)	C/C	The dog does not carry any copies of the tested allele causing curly coat. The dog most likely has non-curly hair.
<i>MC5R</i> c.237A>T	T/T	The dog has two copies of the allele associated with low shedding. The dog is likely average or low shedder.
SGK3 (p.Val96Glyfs)	1/1	The dog does not carry the tested hairlessness allele of the American Hairless Terrier.

On behalf of Genoscoper Laboratories,

boner mas SIGNATURE



 Registered Name:
 Trucker's Freak On A Leash
 Owner:
 Sanna Koponen

 Nickname:
 Bruno
 Country:
 Finland

 Registration ID:
 ER47005/19
 Testing date:
 2019/10/28

 Microchip:
 985141001208561
 DNA
 Identified with standard identification profile:

 Breed:
 Cinneco dell'Etna
 profile:
 JSAG 2006 markers profile:

Dog's identity verified from microchip or tattoo by veterinarian or other authorised person during sample taking: Yes

### Test results - Traits - page 2

#### **Coat Colour**

Trait	Genotype	Description
Colour Locus E - Extensions	e/e	The dog has recessive red coat colour.
Colour Locus B - Brown	b/b	The dog is likely to have brown pigment.
Colour Locus K - Dominant Black	ky/ky	The dog is likely to express the coat colour defined by the colour locus A.
Colour Locus A - Agouti	ay/ay	The dog is genetically sable.
Colour Locus S - Piebald or extreme white spotting	S/S	The dog is likely to have solid coat colour with minimal white.
Colour Locus H - Harlequin	h/h	The dog doesn't have harlequin pattern.
Dilution (d <sup>2</sup> allele)	D/D	The dog does not carry any copies of the rare d2 allele associated with dilution in Chow Chow, French Bulldog, Sloughi and Thai Ridgeback.
Merle (M allele)	m/m	The dog is genetically non-merle and does not carry a <i>SILV</i> gene SINE insertion.
Saddle Tan ( <i>RALY</i> gene dupl.)	-/-	The dog may have saddle tan pattern if it has also tan point genotype at the A locus.
Albinism (caL-allele)	C/C	The dog does not carry the tested mutation for albinism.

On behalf of Genoscoper Laboratories,

mer SIGNATURE



Registered Name:	Trucker's Freak On A Leash	Owner:	Sanna Koponen
Nickname:	Bruno	Country:	Finland
<b>Registration ID:</b>	ER47005/19	Testing date:	2019/10/28
Microchip:	985141001208561		Identified with standard
Breed:	Cirneco dell'Etna	identification profile:	ISAG 2006 markers
Gender:	Male	P	

Dog's identity verified from microchip or tattoo by veterinarian or other authorised person during sample taking: Yes

# Test results - Traits - page 3

### **Body Size**

Trait	Genotype	Description
<i>IGF1</i> (chr15:41221438)	A/G	The dog is heterozygous for the ancestral allele. This means that it carries one copy of the genetic allele typically associated with small body mass and one copy typically associated with large body mass.
<i>IGF1R</i> c.611G>A (p.Arg204His)	A/G	The dog carries one copy of the derived allele and one copy of the ancestral allele.
<i>ACSL4</i> chrX.82919525C>T	C/C	The dog doesn't have the allele associated with large skeletal size and heavy muscling with considerable back fat thickness.
<i>IGSF1</i> p.Asp768Glu	C/C	The dog doesn't have the allele associated with heavy muscling
IRS4 chrX:82296039	G/G	The dog doesn't have the allele associated with large body size.
FGF4 insertion	D/D	The dog is homozygous for the ancient allele. The dog is likely to have legs of normal length.
STC2 (chr4:39182836)	T/T	The dog has two copies of the ancestral allele associated with larger body size.
<i>GHR1</i> (p.Glu191Lys)	G/G	The dog has two copies of the ancestral allele associated with larger body size.
<i>GHR2</i> (p.Pro177Leu)	C/C	The dog has two copies of the ancestral allele associated with larger body size.
<i>HMGA2</i> (chr10:8348804)	A/G	Your dog carries one copy of the derived allele and one copy of the ancestral allele. The dog may have a bit smaller size.

On behalf of Genoscoper Laboratories,

boner onas SIGNATURE



Registered Name:	Trucker's Freak On A Leash	Owner:	Sanna Koponen
Nickname:	Bruno	Country:	Finland
<b>Registration ID:</b>	ER47005/19	Testing date:	2019/10/28
Microchip:	985141001208561		Identified with standard
Breed:	Cirneco dell'Etna	identification profile:	ISAG 2006 markers
Gender:	Male	F	

Dog's identity verified from microchip or tattoo by veterinarian or other authorised person during sample taking: Yes

# Test results - Traits - page 4

### Morphology

Trait	Genotype	Description
<i>BMP3</i> c.1344C>A (p.Phe448Leu)	C/C	The dog does not carry the tested allele typically associated with shortened head (brachycephaly). The dog is more likely to have an elongated head (dolichocephaly).
SMOC2	D/D	The dog does not carry the tested allele typically associated with shortened head (brachycephaly). The dog is more likely to have an elongated head (dolichocephaly).
chr10:11072007	T/T	The dog does not carry an allele typically associated with floppy ears. The dog is more likely to have pricked than floppy ears.
<i>T</i> c.189C>G (p.Ile63Met)	C/C	The dog does not carry the tested bobtail-causing genetic variant. The dog is most likely long-tailed.
<i>EPAS1</i> (p.Gly305Ser)	G/G	The dog does not carry the tested variant associated with adaptation to high altitudes.
LIMBR1 DC-1	G/G	The dog does not carry the tested allele associated with hind dewclaws in Asian breeds. The dog is not likely to have hind dewclaws.
LIMBR1 DC-2	G/G	The dog does not carry the tested allele associated with hind dewclaws in western breeds. The dog is likely not to have hind dewclaws.
AXL4	D/D	The dog does not have the tested allele typically associated with blue eyes in Siberian Huskies. The dog is likely to have brown eyes.

On behalf of Genoscoper Laboratories,

mer onas SIGNATURE



### Blood Disorders - page 1

Bleeding disorder due to P2RY12 defect         Autosomal Recessive         Clear           Canine Cyclic Neutropenia, Cyclic Hematopoiesis, Grey Collie Syndrome, (CN)         Autosomal Recessive         Clear           Canine Leukocyte Adhesion Deficiency (CLAD), type III         Autosomal Recessive         Clear           Canine Scott Syndrome, (CSS)         Autosomal Recessive         Clear           Factor IX Deficiency or Hemophilia B; mutation originally found in Airedale Terrier         X-linked Recessive         Clear           Factor IX Deficiency or Hemophilia B; mutation originally found in Lhasa Apso         X-linked Recessive         Clear           Factor IX Deficiency or Hemophilia A; mutation originally found in Boxer         X-linked Recessive         Clear           Factor VII Deficiency or Hemophilia A; mutation originally found in Boxer         X-linked Recessive         Clear           Factor VIII Deficiency or Hemophilia A; mutation originally found in Old English Sheepdog         X-linked Recessive         Clear           Factor VIII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally found in German Shepherd         X-linked Recessive         Clear           Factor XII Deficiency         Autosomal Recessive         Clear         Clear           Factor XII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally found in German Shepherd         Clear         Clear           Factor XI Deficiency         Autosomal Recessiv	Disorder	Mode of Inheritance	Result
(CN)       Autosomal Recessive       Clear         Canine Leukocyte Adhesion Deficiency (CLAD), type III       Autosomal Recessive       Clear         Canine Scott Syndrome, (CSS)       Autosomal Recessive       Clear         Factor IX Deficiency or Hemophilia B; mutation originally found in Airedale       X-linked Recessive       Clear         Factor IX Deficiency or Hemophilia B; mutation originally found in Lhasa       X-linked Recessive       Clear         Factor IX Deficiency or Hemophilia B; mutation originally found in Lhasa       X-linked Recessive       Clear         Factor IX Deficiency or Hemophilia A; mutation originally found in Boxer       X-linked Recessive       Clear         Factor VII Deficiency or Hemophilia A; mutation originally found in Boxer       X-linked Recessive       Clear         Factor VIII Deficiency or Hemophilia A; mutation originally found in Old       X-linked Recessive       Clear         Factor VIII Deficiency or Hemophilia A; mutation originally found in Old       X-linked Recessive       Clear         Factor VIII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally       X-linked Recessive       Clear         Factor XI Deficiency       Hemophilia A; p.Cys548Tyr mutation originally       X-linked Recessive       Clear         Factor XI Deficiency       Hemophilia A; p.Cys548Tyr mutation originally found in       Autosomal Recessive       Clear         Ganzma	Bleeding disorder due to P2RY12 defect	Autosomal Recessive	Clear
Canine Scott Syndrome, (CSS)Autosomal RecessiveClearFactor IX Deficiency or Hemophilia B; mutation Gly379GluX-linked RecessiveClearFactor IX Deficiency or Hemophilia B; mutation originally found in Airedale TerrierX-linked RecessiveClearFactor IX Deficiency or Hemophilia B; mutation originally found in Lhasa ApsoX-linked RecessiveClearFactor IX Deficiency or Hemophilia B; mutation originally found in Lhasa ApsoX-linked RecessiveClearFactor VII Deficiency or Hemophilia A; mutation originally found in BoxerX-linked RecessiveClearFactor VIII Deficiency or Hemophilia A; mutation originally found in German Shepherd DogX-linked RecessiveClearFactor VIII Deficiency or Hemophilia A; mutation originally found in Old Erator VIII Deficiency or Hemophilia A; mutation originally found in Old In German ShepherdX-linked RecessiveClearFactor VIII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally found in German ShepherdX-linked RecessiveClearFactor XI DeficiencyAutosomal Dominant (Incomplete Penetrance)ClearFamilial Congenital Methemoglobinemia; mutation originally found in Pyrenean Mountain DogAutosomal RecessiveClearGlanzmann Thrombasthenia Type I, (GT); mutation originally found in Pyrenean Mountain DogAutosomal RecessiveClearHereditary ElliptocytosisClearClearClearHereditary Phosphofructokinase (PFK) DeficiencyAutosomal RecessiveClearMacrothrombocytopenia; disease-linked variant originally found in Norfolk and Cairm TerrierAutosomal Recessive		Autosomal Recessive	Clear
Factor IX Deficiency or Hemophilia B; mutation Gly379GluX-linked RecessiveClearFactor IX Deficiency or Hemophilia B; mutation originally found in Airedale TerrierX-linked RecessiveClearFactor IX Deficiency or Hemophilia B; mutation originally found in Lhasa ApsoX-linked RecessiveClearFactor IX Deficiency or Hemophilia B; mutation originally found in Lhasa ApsoX-linked RecessiveClearFactor VII Deficiency or Hemophilia A; mutation originally found in BoxerX-linked RecessiveClearFactor VIII Deficiency or Hemophilia A; mutation originally found in German Shepherd DogX-linked RecessiveClearFactor VIII Deficiency or Hemophilia A; mutation originally found in Old English SheepdogX-linked RecessiveClearFactor VIII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally found in German ShepherdX-linked RecessiveClearFactor XI Deficiencyr Hemophilia A; p.Cys548Tyr mutation originally found in German ShepherdClearClearFactor XI DeficiencyAutosomal Dominant (Incomplete Penetrance)ClearFamilial Congenital Methemoglobinemia; mutation originally found in Pyrenean Mountain DogAutosomal RecessiveClearGlanzmann Thrombasthenia Type I, (GT); mutation originally found in Pyrenean Mountain DogAutosomal RecessiveClearHereditary ElliptocytosisClearClearHereditary Phosphofructokinase (PFK) DeficiencyAutosomal RecessiveClearMacrothrombocytopenia; disease-linked variant originally found in Norfolk and Cairn TerrierAutosomal RecessiveClear<	Canine Leukocyte Adhesion Deficiency (CLAD), type III	Autosomal Recessive	Clear
Factor IX Deficiency or Hemophilia B; mutation originally found in Airedale TerrierX-linked RecessiveClearFactor IX Deficiency or Hemophilia B; mutation originally found in Lhasa ApsoX-linked RecessiveClearFactor VII DeficiencyAutosomal RecessiveClearFactor VIII Deficiency or Hemophilia A; mutation originally found in BoxerX-linked RecessiveClearFactor VIII Deficiency or Hemophilia A; mutation originally found in German Shepherd DogX-linked RecessiveClearFactor VIII Deficiency or Hemophilia A; mutation originally found in OgX-linked RecessiveClearFactor VIII Deficiency or Hemophilia A; mutation originally found in Old English SheepdogX-linked RecessiveClearFactor XI Deficiency or Hemophilia A; p.Cys548Tyr mutation originally found in German ShepherdX-linked RecessiveClearFactor XI DeficiencyAutosomal Dominant (Incomplete Penetrance)ClearFamilal Congenital Methemoglobinemia; mutation originally found in PomeranianAutosomal RecessiveClearGlanzmann Thrombasthenia Type I, (GT); mutation originally found in mixed breed dogsAutosomal RecessiveClearHereditary ElliptocytosisClearClearClearHereditary Phosphofructokinase (PFK) DeficiencyAutosomal RecessiveClearMacrothrombocytopenia; disease-linked variant originally found in Norfolk and Cairn TerrierAutosomal RecessiveClearMay-Hegglin Anomaly (MHA)Autosomal RecessiveClear	Canine Scott Syndrome, (CSS)	Autosomal Recessive	Clear
TerrierClearFactor IX Deficiency or Hemophilia B; mutation originally found in Lhasa ApsoX-linked RecessiveClearFactor VII DeficiencyAutosomal RecessiveClearFactor VIII Deficiency or Hemophilia A; mutation originally found in Boxer German Shepherd DogX-linked RecessiveClearFactor VIII Deficiency or Hemophilia A; mutation originally found in German Shepherd DogX-linked RecessiveClearFactor VIII Deficiency or Hemophilia A; mutation originally found in Old English SheepdogX-linked RecessiveClearFactor VIII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally found in German ShepherdX-linked RecessiveClearFactor XII DeficiencyAutosomal Dominant (Incomplete Penetrance)ClearFactor XI DeficiencyAutosomal Dominant (Incomplete Penetrance)ClearFamilial Congenital Methemoglobinemia; mutation originally found in PomeranianAutosomal RecessiveClearGlanzmann Thrombasthenia Type I, (GT); mutation originally found in mixed breed dogsAutosomal RecessiveClearHereditary ElliptocytosisClearClearClearHereditary Phosphofructokinase (PFK) DeficiencyAutosomal RecessiveClearMacrothrombocytopenia; disease-linked variant originally found in Norfolk and Cairn TerrierAutosomal RecessiveClearMay-Hegglin Anomaly (MHA)Autosomal RecessiveClear	Factor IX Deficiency or Hemophilia B; mutation Gly379Glu	X-linked Recessive	Clear
ApsoAutosomal RecessiveClearFactor VII DeficiencyrecessiveClearFactor VIII Deficiency or Hemophilia A; mutation originally found in BoxerX-linked RecessiveClearFactor VIII Deficiency or Hemophilia A; mutation originally found in German Shepherd DogX-linked RecessiveClearFactor VIII Deficiency or Hemophilia A; mutation originally found in Old English SheepdogX-linked RecessiveClearFactor VIII Deficiency or Hemophilia A; mutation originally found in Old English SheepdogX-linked RecessiveClearFactor VIII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally found in German ShepherdX-linked RecessiveClearFactor XI DeficiencyAutosomal Dominant (Incomplete Penetrance)ClearFactor XI DeficiencyAutosomal RecessiveClearGlanzmann Thrombasthenia Type I, (GT); mutation originally found in Pyrenean Mountain DogAutosomal RecessiveClearGlanzmann Thrombasthenia Type I, (GT); mutation originally found in mixed breed dogsAutosomal RecessiveClearHereditary PliptocytosisClearClearHereditary Phosphofructokinase (PFK) DeficiencyAutosomal RecessiveClearMacrothrombocytopenia; disease-linked variant originally found in Norfolk and Cairm TerrierAutosomal RecessiveClearMay-Hegglin Anomaly (MHA)Autosomal Dominant Autosomal DominantClear		X-linked Recessive	Clear
Factor VIII Deficiency or Hemophilia A; mutation originally found in BoxerX-linked RecessiveClearFactor VIII Deficiency or Hemophilia A; mutation originally found in German Shepherd DogX-linked RecessiveClearFactor VIII Deficiency or Hemophilia A; mutation originally found in Old English SheepdogX-linked RecessiveClearFactor VIII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally found in German ShepherdX-linked RecessiveClearFactor XI DeficiencyAutosomal Dominant (Incomplete Penetrance)ClearFamilial Congenital Methemoglobinemia; mutation originally found in PomeranianAutosomal RecessiveClearGlanzmann Thrombasthenia Type I, (GT); mutation originally found in mixed breed dogsAutosomal RecessiveClearHereditary ElliptocytosisClearClearMacrothrombocytopenia; disease-linked variant originally found in Norfolk and Cairn TerrierAutosomal RecessiveClearMay-Hegglin Anomaly (MHA)Autosomal RecessiveClear		X-linked Recessive	Clear
Factor VIII Deficiency or Hemophilia A; mutation originally found in German Shepherd DogX-linked RecessiveClearFactor VIII Deficiency or Hemophilia A; mutation originally found in Old English SheepdogX-linked RecessiveClearFactor VIII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally found in German ShepherdX-linked RecessiveClearFactor VIII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally found in German ShepherdX-linked RecessiveClearFactor XI DeficiencyAutosomal Dominant (Incomplete Penetrance)ClearFamilial Congenital Methemoglobinemia; mutation originally found in PomeranianAutosomal RecessiveClearGlanzmann Thrombasthenia Type I, (GT); mutation originally found in mixed breed dogsAutosomal RecessiveClearHereditary ElliptocytosisClearClearClearHereditary Phosphofructokinase (PFK) DeficiencyAutosomal RecessiveClearMacrothrombocytopenia; disease-linked variant originally found in Norfolk and Cairn TerrierAutosomal RecessiveClearMay-Hegglin Anomaly (MHA)Autosomal Dominant clearClear	Factor VII Deficiency	Autosomal Recessive	Clear
German Shepherd DogClearFactor VIII Deficiency or Hemophilia A; mutation originally found in Old English SheepdogX-linked RecessiveClearFactor VIII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally found in German ShepherdX-linked RecessiveClearFactor XI DeficiencyAutosomal Dominant (Incomplete Penetrance)ClearFactor XI DeficiencyAutosomal Dominant (Incomplete Penetrance)ClearFamilial Congenital Methemoglobinemia; mutation originally found in PomeranianAutosomal RecessiveClearGlanzmann Thrombasthenia Type I, (GT); mutation originally found in Pyrenean Mountain DogAutosomal RecessiveClearGlanzmann Thrombasthenia Type I, (GT); mutation originally found in mixed breed dogsAutosomal RecessiveClearHereditary ElliptocytosisClearMacrothrombocytopenia; disease-linked variant originally found in Norfolk and Caim TerrierAutosomal RecessiveClearMay-Hegglin Anomaly (MHA)Autosomal DominantClear	Factor VIII Deficiency or Hemophilia A; mutation originally found in Boxer	X-linked Recessive	Clear
English SheepdogClearFactor VIII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally found in German ShepherdX-linked RecessiveClearFactor XI DeficiencyAutosomal Dominant (Incomplete Penetrance)ClearFamilial Congenital Methemoglobinemia; mutation originally found in PomeranianAutosomal RecessiveClearGlanzmann Thrombasthenia Type I, (GT); mutation originally found in Pyrenean Mountain DogAutosomal RecessiveClearGlanzmann Thrombasthenia Type I, (GT); mutation originally found in mixed breed dogsAutosomal RecessiveClearHereditary ElliptocytosisClearClearMacrothrombocytopenia; disease-linked variant originally found in Norfolk and Caim TerrierAutosomal RecessiveClearMay-Hegglin Anomaly (MHA)Autosomal Norfolk Autosomal DominantClear		X-linked Recessive	Clear
found in German ShepherdAutosomal Dominant (Incomplete Penetrance)ClearFactor XI DeficiencyAutosomal Dominant (Incomplete Penetrance)ClearFamilial Congenital Methemoglobinemia; mutation originally found in PomeranianAutosomal RecessiveClearGlanzmann Thrombasthenia Type I, (GT); mutation originally found in Pyrenean Mountain DogAutosomal RecessiveClearGlanzmann Thrombasthenia Type I, (GT); mutation originally found in mixed breed dogsAutosomal RecessiveClearHereditary ElliptocytosisClearClearHereditary Phosphofructokinase (PFK) DeficiencyAutosomal RecessiveClearMacrothrombocytopenia; disease-linked variant originally found in Norfolk and Cairn TerrierAutosomal RecessiveClearMay-Hegglin Anomaly (MHA)Autosomal DominantClear		X-linked Recessive	Clear
Familial Congenital Methemoglobinemia; mutation originally found in PomeranianAutosomal RecessiveClearGlanzmann Thrombasthenia Type I, (GT); mutation originally found in Pyrenean Mountain DogAutosomal RecessiveClearGlanzmann Thrombasthenia Type I, (GT); mutation originally found in mixed breed dogsAutosomal RecessiveClearGlanzmann Thrombasthenia Type I, (GT); mutation originally found in mixed breed dogsAutosomal RecessiveClearHereditary ElliptocytosisClearClearHereditary Phosphofructokinase (PFK) DeficiencyAutosomal RecessiveClearMacrothrombocytopenia; disease-linked variant originally found in Norfolk and Caim TerrierAutosomal RecessiveClearMay-Hegglin Anomaly (MHA)Autosomal DominantClear		X-linked Recessive	Clear
Pomeranian       Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Pyrenean Mountain Dog       Autosomal Recessive       Clear         Glanzmann Thrombasthenia Type I, (GT); mutation originally found in mixed breed dogs       Autosomal Recessive       Clear         Hereditary Elliptocytosis       Clear       Clear         Hereditary Phosphofructokinase (PFK) Deficiency       Autosomal Recessive       Clear         Macrothrombocytopenia; disease-linked variant originally found in Norfolk       Autosomal Recessive       Clear         May-Hegglin Anomaly (MHA)       Autosomal Dominant       Clear	Factor XI Deficiency		Clear
Pyrenean Mountain Dog       Glanzmann Thrombasthenia Type I, (GT); mutation originally found in mixed breed dogs       Autosomal Recessive       Clear         Hereditary Elliptocytosis       Clear         Hereditary Phosphofructokinase (PFK) Deficiency       Autosomal Recessive       Clear         Macrothrombocytopenia; disease-linked variant originally found in Norfolk and Cairn Terrier       Autosomal Recessive       Clear         May-Hegglin Anomaly (MHA)       Autosomal Dominant       Clear		Autosomal Recessive	Clear
mixed breed dogs     Clear       Hereditary Elliptocytosis     Clear       Hereditary Phosphofructokinase (PFK) Deficiency     Autosomal Recessive     Clear       Macrothrombocytopenia; disease-linked variant originally found in Norfolk and Cairm Terrier     Autosomal Recessive     Clear       May-Hegglin Anomaly (MHA)     Autosomal Dominant     Clear		Autosomal Recessive	Clear
Hereditary Phosphofructokinase (PFK) DeficiencyAutosomal RecessiveClearMacrothrombocytopenia; disease-linked variant originally found in Norfolk and Cairn TerrierAutosomal RecessiveClearMay-Hegglin Anomaly (MHA)Autosomal DominantClear		Autosomal Recessive	Clear
Macrothrombocytopenia; disease-linked variant originally found in Norfolk and Cairn Terrier       Autosomal Recessive       Clear         May-Hegglin Anomaly (MHA)       Autosomal Dominant       Clear	Hereditary Elliptocytosis		Clear
and Cairn Terrier Autosomal Dominant Clear	Hereditary Phosphofructokinase (PFK) Deficiency	Autosomal Recessive	Clear
	Macrothrombocytopenia; disease-linked variant originally found in Norfolk and Cairn Terrier	Autosomal Recessive	Clear
Prekallikrein Deficiency Autosomal Recessive Clear	May-Hegglin Anomaly (MHA)	Autosomal Dominant	Clear
	Prekallikrein Deficiency	Autosomal Recessive	Clear



### Blood Disorders - page 2

Disorder	Mode of Inheritance	Result
Pyruvate Kinase Deficiency; mutation originally found in Basenji	Autosomal Recessive	Clear
Pyruvate Kinase Deficiency; mutation originally found in Beagle	Autosomal Recessive	Clear
Pyruvate Kinase Deficiency; mutation originally found in Pug	Autosomal Recessive	Clear
Pyruvate Kinase Deficiency; mutation originally found in West Highland White Terrier	Autosomal Recessive	Clear
Thrombopathia; mutation originally found in Basset Hound	Autosomal Recessive	Clear
Thrombopathia; mutation originally found in Eskimo Spitz	Autosomal Recessive	Clear
Thrombopathia; mutation originally found in Landseer	Autosomal Recessive	Clear
Trapped Neutrophil Syndrome, (TNS)	Autosomal Recessive	Clear
Von Willebrand's Disease (vWD) Type 1	Autosomal Recessive	Clear
Von Willebrand's Disease (vWD) Type 2	Autosomal Recessive	Clear
Von Willebrand's Disease (vWD) Type 3; mutation originally found in Kooikerhondje	Autosomal Recessive	Clear
Von Willebrand's Disease (vWD) Type 3; mutation originally found in Scottish Terrier	Autosomal Recessive	Clear
Von Willebrand's Disease (vWD) Type 3; mutation originally found in Shetland Sheepdog	Autosomal Recessive	Clear



### Ocular Disorders - page 1

Disorder	Mode of Inheritance	Result
Canine Multifocal Retinopathy 1, (CMR1); mutation originally found in Mastiff-related breeds	Autosomal Recessive	Clear
Canine Multifocal Retinopathy 2, (CMR2); mutation originally found in Coton de Tulear	Autosomal Recessive	Clear
Canine Multifocal Retinopathy 3, (CMR3); mutation originally found in Lapponian Herder	Autosomal Recessive	Clear
Cone Degeneration, (CD) or Achromatopsia; mutation originally found in Alaskan Malamute	Autosomal Recessive	Clear
Cone Degeneration, (CD) or Achromatopsia; mutation originally found in German Shepherd Dog	Autosomal Recessive	Clear
Cone Degeneration, (CD) or Achromatopsia; mutation originally found in German Shorthaired Pointer	Autosomal Recessive	Clear
Cone-Rod Dystrophy 1, (crd1); mutation originally found in American Staffordshire Terrier	Autosomal Recessive	Clear
Cone-Rod Dystrophy 2, (crd2); mutation originally found in American Pit Bull Terrier	Autosomal Recessive	Clear
Cone-Rod Dystrophy, (cord1-PRA / crd4)	Autosomal Recessive (Incomplete Penetrance)	Clear
Cone-Rod Dystrophy, Standard Wirehaired Dachshund, (crd SWD)	Autosomal Recessive	Clear
Congenital Eye Disease; mutation originally found in Irish Soft-Coated Wheaten Terrier	Autosomal Recessive	Clear
Dominant Progressive Retinal Atrophy, (DPRA)	Autosomal Dominant	Clear
Early Retinal Degeneration, (erd); mutation originally found in Norwegian Elkhound	Autosomal Recessive	Clear
Generalized Progressive Retinal Atrophy	Autosomal Recessive	Clear
Golden Retriever Progressive Retinal Atrophy 1, (GR_PRA 1)	Autosomal Recessive	Clear
Golden Retriever Progressive Retinal Atrophy 2, (GR_PRA 2)	Autosomal Recessive	Clear
Primary Hereditary Cataract, (PHC); mutation originally found in Australian Shepherd	Autosomal Dominant (Incomplete Penetrance)	Clear
Primary Lens Luxation, (PLL)	Autosomal Recessive	Clear
Primary Open Angle Glaucoma, (POAG); mutation originally found in Basset Fauve de Bretagne	Autosomal Recessive	Clear
Primary Open Angle Glaucoma, (POAG); mutation originally found in Beagle	Autosomal Recessive	Clear
Primary Open Angle Glaucoma, (POAG); mutation originally found in Norwegian Elkhound	Autosomal Recessive	Clear



### Ocular Disorders - page 2

Disorder	Mode of Inheritance	Result
Primary Open Angle Glaucoma, (POAG); mutation originally found in Petit Basset Griffon Vendeen	Autosomal Recessive	Clear
Primary lens luxation (PLL) and glaucoma; mutation originally found in Shar Pei	Autosomal Recessive	Clear
Progressive Retinal Atrophy (PRA4); mutation originally found in Lhasa Apso	Autosomal Recessive	Clear
Progressive Retinal Atrophy Type III, (PRA type III); mutation originally found in Tibetan Spaniel and Tibetan Terrier	Autosomal Recessive	Clear
Progressive Retinal Atrophy, (CNGA1-PRA); mutation originally found in Shetland Sheepdog	Autosomal Recessive	Clear
Progressive Retinal Atrophy, (PAP1_PRA); mutation originally found in Papillon and Phalene	Autosomal Recessive	Clear
Progressive Retinal Atrophy, (PRA); mutation originally found in Basenji	Autosomal Recessive	Clear
Progressive Retinal Atrophy, (PRA); mutation originally found in Swedish Vallhund	Autosomal Recessive	Clear
Rod-Cone Dysplasia 1, (rcd1); mutation originally found in Irish Setter	Autosomal Recessive	Clear
Rod-Cone Dysplasia 1a, (rdc1a); mutation originally found in Sloughi	Autosomal Recessive	Clear
Rod-Cone Dysplasia 3, (rcd3)	Autosomal Recessive	Clear
X-Linked Progressive Retinal Atrophy 1, (XLPRA1)	X-linked Recessive	Clear
X-Linked Progressive Retinal Atrophy 2, (XLPRA2; Type A PRA)	X-linked Recessive	Clear

#### **Cardiac Disorders**

Disorder	Mode of Inheritance	Result
Dilated Cardiomyopathy, (DCM); mutation originally found in Schnauzer	Autosomal Recessive	Clear
Long QT Syndrome	Autosomal Dominant	Clear



#### **Endocrine Disorders**

Disorder	Mode of Inheritance	Result
Congenital Hypothyroidism; mutation originally found in Tenterfield Terrier	Autosomal Recessive	Clear
Congenital Hypothyroidism; mutation originally found in Toy Fox and Rat Terrier	Autosomal Recessive	Clear

### Immunological Disorders

Disorder	Mode of Inheritance	Result
Autosomal Recessive Severe Combined Immunodeficiency, (ARSCID)	Autosomal Recessive	Clear
Complement 3 (C3) Deficiency	Autosomal Recessive	Clear
Myeloperoxidase Deficiency	Autosomal Recessive	Clear
Severe Combined Immunodeficiency in Frisian Water Dogs, (SCID)	Autosomal Recessive	Clear
X-Linked Severe Combined Immunodeficiency (XSCID); mutation originally found in Basset Hound	X-linked Recessive	Clear
X-Linked Severe Combined Immunodeficiency (XSCID); mutation originally found in Cardigan Welsh Corgi	X-linked Recessive	Clear



#### **Renal Disorders**

Disorder	Mode of Inheritance	Result
2,8-Dihydroxyadenine (2,8-DHA) urolithiasis	Autosomal Recessive	Clear
Cystinuria Type I-A; mutation originally found in Newfoundland Dog	Autosomal Recessive	Clear
Cystinuria Type II-A; mutation originally found in Australian Cattle Dog	Autosomal Dominant	Clear
Familial Nephropathy (FN); mutation originally found in English Cocker Spaniel	Autosomal Recessive	Clear
Familial Nephropathy (FN); mutation originally found in English Springer Spaniel	Autosomal Recessive	Clear
Fanconi Syndrome	Autosomal Recessive	Clear
Hyperuricosuria, (HUU)	Autosomal Recessive	Clear
Polycystic Kidney Disease in Bull Terriers, (BTPKD)	Autosomal Dominant	Clear
Primary Hyperoxaluria, (PH); mutation originally found in Coton de Tulear	Autosomal Recessive	Clear
Protein Losing Nephropathy, (PLN); NPHS1 gene variant		Clear
Renal Cystadenocarcinoma and Nodular Dermatofibrosis, (RCND)	Autosomal Dominant	Clear
X-Linked Hereditary Nephropathy, (XLHN)	X-linked Recessive	Clear
X-Linked Hereditary Nephropathy, (XLHN); mutation originally found in Navasota Dog	X-linked Recessive	Clear
Xanthinuria, Type 1a; mutation originally found in mixed breed dogs	Autosomal Recessive	Clear
Xanthinuria, Type 2a; mutation originally found in Toy Manchester Terrier	Autosomal Recessive	Clear
Xanthinuria, Type 2b; mutation originally found in Cavalier King Charles Spaniel and English Cocker Spaniel	Autosomal Recessive	Clear



#### **Metabolic Disorders**

Disorder	Mode of Inheritance	Result
Glycogen Storage Disease Type II or Pompe's Disease, (GSD II)	Autosomal Recessive	Clear
Glycogen Storage Disease Type IIIa, (GSD IIIa)	Autosomal Recessive	Clear
Glycogen Storage Disease Type Ia, (GSD Ia)	Autosomal Recessive	Clear
Hypocatalasia or Acatalasemia	Autosomal Recessive	Clear
Intestinal Cobalamin Malabsorption or Imerslund-Gräsbeck Syndrome, (IGS); mutation originally found in Beagle	Autosomal Recessive	Clear
Intestinal Cobalamin Malabsorption or Imerslund-Gräsbeck Syndrome, (IGS); mutation originally found in Border Collie	Autosomal Recessive	Clear
Mucopolysaccharidosis Type IIIA, (MPS IIIA); mutation originally found in Dachshund	Autosomal Recessive	Clear
Mucopolysaccharidosis Type IIIA, (MPS IIIA); mutation originally found in New Zealand Huntaway	Autosomal Recessive	Clear
Mucopolysaccharidosis Type VII, (MPS VII); mutation originally found in Brazilian Terrier	Autosomal Recessive	Clear
Mucopolysaccharidosis Type VII, (MPS VII); mutation originally found in German Shepherd	Autosomal Recessive	Clear
Pyruvate Dehydrogenase Phosphatase 1 (PDP1) Deficiency	Autosomal Recessive	Clear



#### **Muscular Disorders**

Mode of Inheritance	Result
X-linked Recessive	Clear
Autosomal Recessive	Clear
Autosomal Recessive	Clear
X-linked Recessive	Clear
X-linked Recessive	Clear
Autosomal Recessive	Clear
Autosomal Recessive	Clear
Autosomal Recessive	Clear
Autosomal Recessive	Clear
X-linked Recessive	Clear
Autosomal Recessive	Clear
X-linked Recessive	Clear
	X-linked Recessive Autosomal Recessive Autosomal Recessive X-linked Recessive X-linked Recessive Autosomal Recessive Autosomal Recessive Autosomal Recessive Autosomal Recessive X-linked Recessive Autosomal Recessive



### Neurological Disorders - page 1

Disorder	Mode of Inheritance	Result
Acral Mutilation Syndrome, (AMS)	Autosomal Recessive	Clear
Alaskan Husky Encephalopathy, (AHE)	Autosomal Recessive	Clear
Alexander Disease (AxD); mutation originally found in Labrador Retriever	Autosomal Dominant	Clear
Bandera's Neonatal Ataxia, (BNAt)	Autosomal Recessive	Clear
Benign Familial Juvenile Epilepsy or Remitting Focal Epilepsy	Autosomal Recessive	Clear
Cerebellar Cortical Degeneration, (CCD); mutation originally found in Vizsla	Autosomal Recessive	Clear
Cerebral Dysfunction; mutation originally found in Friesian Stabyhoun	Autosomal Recessive	Clear
Dandy-Walker-Like Malformation (DWLM); mutation originally found in Eurasier	Autosomal Recessive	Clear
Degenerative Myelopathy, (DM; SOD1A)	Autosomal Recessive (Incomplete Penetrance)	Clear
Early-Onset Progressive Polyneuropathy; mutation originally found in Alaskan Malamute	Autosomal Recessive	Clear
Early-Onset Progressive Polyneuropathy; mutation originally found in Greyhound	Autosomal Recessive	Clear
Fetal Onset Neuroaxonal Dystrophy, (FNAD)	Autosomal Recessive	Clear
Hereditary Ataxia or Cerebellar Ataxia; mutation originally found in Old English Sheepdog and Gordon Setter	Autosomal Recessive	Clear
Hereditary Ataxia; mutation originally found in in Norwegian Buhund	Autosomal Recessive	Clear
Hyperekplexia or Startle Disease	Autosomal Recessive	Clear
Hypomyelination; mutation originally found in Weimaraner	Autosomal Recessive	Clear
Juvenile Myoclonic Epilepsy, (JME); mutation originally found in Rhodesian Ridgeback	Autosomal Recessive	Clear
Juvenile encephalopathy; mutation originally found in Parson Russell Terrier	Autosomal Recessive	Clear
L-2-Hydroxyglutaric aciduria, (L2HGA); mutation originally found in Staffordshire Bull Terrier	Autosomal Recessive	Clear
L-2-Hydroxyglutaric aciduria, (L2HGA); mutation originally found in West Highland White Terrier	Autosomal Recessive	Clear
Lagotto Storage Disease, (LSD)	Autosomal Recessive	Clear



### Neurological Disorders - page 2

Disorder	Mode of Inheritance	Result
Neonatal Cerebellar Cortical Degeneration or Cerebellar Abiotrophy, (NCCD)	Autosomal Recessive	Clear
Neonatal Encephalopathy with Seizures, (NEWS)	Autosomal Recessive	Clear
Neuroaxonal Dystrophy (NAD); mutation originally found in Spanish Water Dog	Autosomal Recessive	Clear
Neuroaxonal Dystrophy, (NAD); mutation originally found in Papillon	Autosomal Recessive	Clear
Neuronal Ceroid Lipofuscinosis 1, (NCL1); mutation originally found in Dachshund	Autosomal Recessive	Clear
Neuronal Ceroid Lipofuscinosis 10, (NCL10); mutation originally found in American Bulldog	Autosomal Recessive	Clear
Neuronal Ceroid Lipofuscinosis 4A, (NCL4); mutation originally found in American Staffordshire Terrier	Autosomal Recessive	Clear
Neuronal Ceroid Lipofuscinosis 5, (NCL5); mutation originally found in Border Collie	Autosomal Recessive	Clear
Neuronal Ceroid Lipofuscinosis 8, (NCL8); mutation originally found in Alpine Dachsbracke	Autosomal Recessive	Clear
Neuronal Ceroid Lipofuscinosis 8, (NCL8); mutation originally found in Australian Shepherd	Autosomal Recessive	Clear
Neuronal Ceroid Lipofuscinosis 8, (NCL8); mutation originally found in English Setter	Autosomal Recessive	Clear
Neuronal Ceroid Lipofuscinosis, (NCL7); mutation originally found in Chinese Crested Dog and Chihuahua	Autosomal Recessive	Clear
Polyneuropathy with ocular abnormalities and neuronal vacuolation, (POANV); mutation originally found in Black Russian Terrier	Autosomal Recessive	Clear
Progressive Early-Onset Cerebellar Ataxia; mutation originally found in Finnish Hound	Autosomal Recessive	Clear
Sensory Neuropathy; mutation originally found in Border Collie	Autosomal Recessive	Clear
Shaking Puppy Spongiform LeucoEncephaloMyelopathy, (SLEM); mutation originally found in Border Terrier	Autosomal Recessive	Clear
Spinocerebellar Ataxia with Myokymia and/or Seizures (SCA)	Autosomal Recessive	Clear
Spinocerebellar Ataxia/ Late-Onset Ataxia (SCA, LOA)	Autosomal Recessive	Clear
Spongy Degeneration with Cerebellar Ataxia, (SDCA1); mutation originally found in Belgian Shepherd Dog	Autosomal Recessive	Clear
Spongy Degeneration with Cerebellar Ataxia, (SDCA2); mutation originally found in Belgian Shepherd Dog	Autosomal Recessive	Clear
X-Linked Tremors; mutation originally found in English Springer Spaniel	X-linked Recessive	Clear



#### **Neuromuscular Disorders**

Disorder	Mode of Inheritance	Result
Congenital Myasthenic Syndrome (CMS); mutation originally found in Labrador Retriever	Autosomal Recessive	Clear
Congenital Myasthenic Syndrome, (CMS); mutation originally found in Jack Russell Terrier	Autosomal Recessive	Clear
Congenital Myasthenic Syndrome, (CMS); mutation originally found in Old Danish Pointing Dog	Autosomal Recessive	Clear
Exercise-Induced Collapse, (EIC)	Autosomal Recessive (Incomplete Penetrance)	Clear
GM1 Gangliosidosis; mutation originally found in Alaskan Husky	Autosomal Recessive	Clear
GM1 Gangliosidosis; mutation originally found in Portuguese Water Dog	Autosomal Recessive	Clear
GM1 Gangliosidosis; mutation originally found in Shiba Dog	Autosomal Recessive	Clear
GM2 Gangliosidosis, mutation originally found in Japanese Chin	Autosomal Recessive	Clear
GM2 Gangliosidosis; mutation originally found in Toy Poodle	Autosomal Recessive	Clear
Globoid Cell Leukodystrophy or Krabbe Disease, (GLD); mutation originally found in Irish Setter	Autosomal Recessive	Clear
Globoid Cell Leukodystrophy or Krabbe Disease, (GLD); mutation originally found in Terriers	Autosomal Recessive	Clear
Paroxysmal Dyskinesia, (PxD); mutation originally found in Irish Soft Coated Wheaten Terrier	Autosomal Recessive	Clear



### **Skeletal Disorders**

	Result
al Recessive	Clear
al Recessive	Clear
al Recessive	Clear
al Dominant ete Penetrance)	Clear
al Recessive	Clear
al Recessive	Clear
al Recessive	Clear
al Dominant	Clear
al Dominant	Clear
al Recessive	Clear
a	I Recessive



#### **Dermal Disorders**

Disorder	Mode of Inheritance	Result
Dystrophic Epidermolysis Bullosa; mutation originally found in Central Asian Ovcharka	Autosomal Recessive	Clear
Dystrophic Epidermolysis Bullosa; mutation originally found in Golden Retriever	Autosomal Recessive	Clear
Epidermolytic Hyperkeratosis	Autosomal Recessive	Clear
Focal Non-Epidermolytic Palmoplantar Keratoderma, (FNEPPK); mutation originally found in Dogue de Bordeaux	Autosomal Recessive	Clear
Golden Retriever Ichthyosis	Autosomal Recessive	Clear
Hereditary Footpad Hyperkeratosis, (HFH)	Autosomal Recessive	Clear
Hereditary Nasal Parakeratosis, (HNPK); mutation originally found in Greyhound	Autosomal Recessive	Clear
Ichthyosis; mutation originally found in American Bulldog	Autosomal Recessive	Clear
Ichthyosis; mutation originally found in Great Dane	Autosomal Recessive	Clear
Lamellar Ichthyosis, (LI)	Autosomal Recessive	Clear
Lethal Acrodermatitis, (LAD); mutation originally found in in Bull Terrier and Miniature Bull Terrier	Autosomal Recessive	Clear
Ligneous Membranitis	Autosomal Recessive	Clear
Musladin-Lueke syndrome, (MLS)	Autosomal Recessive	Clear
X-Linked Ectodermal Dysplasia, (XHED)	X-linked Recessive	Clear



#### **Other Disorders**

Disorder	Mode of Inheritance	Result
Acute Respiratory Distress Syndrome, (ARDS); mutation originally found in Dalmatian	Autosomal Recessive	Clear
Amelogenesis Imperfecta, (AI); mutation originally found in Italian Greyhound	Autosomal Recessive	Clear
Amelogenesis Imperfecta, (AI); mutation originally found in Parson Russell Terrier	Autosomal Recessive	Clear
Congenital Keratoconjunctivitis Sicca and Ichthyosiform Dermatosis, (CKCSID)	Autosomal Recessive	Clear
Dental Hypomineralisation; mutation originally found in Border Collie	Autosomal Recessive	Clear
Lung Developmental Disease; mutation originally found in in Airedale Terrier	Autosomal Recessive	Clear
Narcolepsy; mutation originally found in Dachshund	Autosomal Recessive	Clear
Narcolepsy; mutation originally found in Doberman Pinscher	Autosomal Recessive	Clear
Narcolepsy; mutation originally found in Labrador Retriever	Autosomal Recessive	Clear
Persistent Müllerian Duct Syndrome, (PMDS); mutation originally found in Miniature Schnauzer	Autosomal Recessive	Clear
Primary Ciliary Dyskinesia, (PCD)	Autosomal Recessive	Clear



### APPENDIX Explanation of the results of the tested disorders

Autosomal recessive inheritance (ARI)

Clear - A dog carries no copies of the tested mutation and has no or reduced likelihood of developing and passing on the disease/condition.

Carrier - A dog carries one copy of the tested mutation. Carriers typically have a normal, healthy appearance but pass on the mutation to approximately 50% of their offspring.

At risk - A dog carries two copies of the tested mutation and is at high or increased risk of developing the disease/condition.

Autosomal dominant inheritance (ADI)

Clear - A dog carries no copies of the tested mutation and has no or reduced likelihood of developing and passing on the disease/condition.

At risk - A dog carries one or two copies of the tested mutation and is at high or increased risk of developing the disease/condition.

X-linked recessive inheritance (X-linked)

Clear - A dog carries no copies of the tested mutation and has no or reduced likelihood of developing and passing on the disease/condition.

Carrier - Female carriers typically have a normal, healthy appearance but carry one copy of the tested mutation on one of their X chromosomes. As males only have one X chromosome, there are no male carriers.

At risk - Female dogs at risk carry two mutated copies of the tested mutation. Males carry one copy of the tested mutation on their single X chromosome. Dogs at risk are at high or increased risk of developing the disease/condition.

Please note that the descriptions above are generalized based on typically observed inheritance patterns. When obtaining a 'carrier' or 'at risk' test result, always refer to the corresponding online test documentation for more detailed information on the condition and any exceptions.

### Genoscoper Laboratories - Legal Notice

Genoscoper Laboratories' services and test results are produced based on samples and materials supplied by the Client. Testing and analysis is performed by using methods and processes that Genoscoper Laboratories deems appropriate. Genoscoper Laboratories reserves the right to make changes in the collection of the single-gene tests included in the testing service as well as to remove results derived from them, if new information comes available that in any way questions the validity of the test results. Results provided by Genoscoper Laboratories are prepared solely for the use of the Client. For further information, please visit: <u>www.mydogdna.com/legal-notices</u>

Registered Name: Trucker's Freak On A Leash 15/8/2020 ID 8700 3696 3300 Nickname: Bruno Registration ID: ER47005/19 Microchip: 985141001208561 Breed: Cirneco dell'Etna Gender: Male Owner: Sanna Koponen Country: Finland Testing date: 28/10/2019 DNA identification profile: Identified with standard ISAG 2006 markers Dog's identity verified from microchip or tattoo by veterinarian or other authorised person during sample taking: Yes **Certificate of DNA Identification Profile**  
 INU055
 INU030
 INU005
 FH2848
 AMELOGENIN
 AHT137
 REN54P11

 210/218
 144/150
 126/126
 230/240
 Y/X
 147/147
 228/232
 REN64E19 REN247M23 REN169O18 REN169D01 INU055 INU030 145/149 266/268 162/170 216/218 147/147 228/232 
 CXX279
 AHTK253
 AHTH260
 AHTH171
 AHTH130

 118/126
 288/292
 248/250
 219/233
 119/119
 REN162C04 REN105LO3 INRA21 FH2054 AHTK211 AHT121 200/204 237/241 99/103 168/168 87/91 102/104 On behalf of Genoscoper Laboratories, fonas Domeir SIGNATURE PROVIDED BY GENOSCOPER LABORATORIES Jonas Donner, PhD, Head of Research and Development at Genoscoper Laboratories PO Box. 1040 FI-00251 Helsinki | info@genoscoper.com