



Nickname: Paavo Country: Finland

Registration ID: ER47004/19 Testing date: 2019/10/28

Microchip: 985141001209254

Breed: Cirneco dell'Etna

DNA Identified with standard identification ISAG 2006 markers profile:

Gender: Male

Dog's identity verified from microchip or tattoo by veterinarian or other authorised person during sample taking: Yes

Test results for pharmacogenetics

Disorder	Mode of Inheritance	Result
Multi-Drug Resistance 1, (MDR1)	Autosomal Dominant	Clear

On behalf of Genoscoper Laboratories,

SIGNATURE





Nickname: Paavo Country: Finland

Registration ID: ER47004/19 Testing date: 2019/10/28

Microchip: 985141001209254

Breed: Cirneco dell'Etna

DNA Identified with standard identification ISAG 2006 markers profile:

Gender: Male

Dog's identity verified from microchip or tattoo by veterinarian or other authorised person during sample taking: Yes

Test results - Traits - page 1

Coat Type

Genotype	Description
L/L	The dog is likely to have short-haired coat.
GG/CC	The dog is not genetically likely to express furnishings.
C/C	The dog does not carry any copies of the tested allele causing curly coat. The dog most likely has non-curly hair.
T/T	The dog has two copies of the allele associated with low shedding. The dog is likely average or low shedder.
1/1	The dog does not carry the tested hairlessness allele of the American Hairless Terrier.
	L/L GG/CC C/C T/T

On behalf of Genoscoper Laboratories,

SIGNATURE





Nickname: Paavo Country: Finland

Registration ID: ER47004/19 Testing date: 2019/10/28

Microchip: 985141001209254

Breed: Cirneco dell'Etna

DNA Identified with standard identification ISAG 2006 markers profile:

Gender: Male

Dog's identity verified from microchip or tattoo by veterinarian or other authorised person during sample taking: Yes

Test results - Traits - page 2

Coat Colour

Trait	Genotype	Description
Colour Locus E - Extensions	e/e	The dog has recessive red coat colour.
Colour Locus B - Brown	b/b	The dog is likely to have brown pigment.
Colour Locus K - Dominant Black	ky/ky	The dog is likely to express the coat colour defined by the colour locus A.
Colour Locus A - Agouti	ay/aw	The dog is genetically sable. The dog carries wolf gray coat colour.
Colour Locus S - Piebald or extreme white spotting	S/S	The dog is likely to have solid coat colour with minimal white.
Colour Locus H - Harlequin	h/h	The dog doesn't have harlequin pattern.
Dilution (d ² allele)	D/D	The dog does not carry any copies of the rare d2 allele associated with dilution in Chow Chow, French Bulldog, Sloughi and Thai Ridgeback.
Merle (M allele)	m/m	The dog is genetically non-merle and does not carry a <i>SILV</i> gene SINE insertion.
Saddle Tan (RALY gene dupl.)	-/dup	The dog may have saddle tan pattern if it has also tan point genotype at the A locus.
Albinism (caL-allele)	C/C	The dog does not carry the tested mutation for albinism.
· · · · · · · · · · · · · · · · · · ·		

On behalf of Genoscoper Laboratories,

SIGNATURE





Nickname: Paavo Country: Finland

Registration ID: ER47004/19 Testing date: 2019/10/28

Microchip: 985141001209254

Breed: Cirneco dell'Etna

DNA Identified with standard identification ISAG 2006 markers profile:

Gender: Male

Dog's identity verified from microchip or tattoo by veterinarian or other authorised person during sample taking: Yes

Test results - Traits - page 3

Body Size

Trait	Genotype	Description
IGF1 (chr15:41221438)	A/G	The dog is heterozygous for the ancestral allele. This means that it carries one copy of the genetic allele typically associated with small body mass and one copy typically associated with large body mass.
IGF1R c.611G>A (p.Arg204His)	A/G	The dog carries one copy of the derived allele and one copy of the ancestral allele.
ACSL4 chrX.82919525C>T	C/C	The dog doesn't have the allele associated with large skeletal size and heavy muscling with considerable back fat thickness.
IGSF1 p.Asp768Glu	C/C	The dog doesn't have the allele associated with heavy muscling
IRS4 chrX:82296039	G/G	The dog doesn't have the allele associated with large body size.
FGF4 insertion	D/D	The dog is homozygous for the ancient allele. The dog is likely to have legs of normal length.
STC2 (chr4:39182836)	T/T	The dog has two copies of the ancestral allele associated with larger body size.
GHR1 (p.Glu191Lys)	G/G	The dog has two copies of the ancestral allele associated with larger body size.
GHR2 (p.Pro177Leu)	C/C	The dog has two copies of the ancestral allele associated with larger body size.
HMGA2 (chr10:8348804)	A/G	Your dog carries one copy of the derived allele and one copy of the ancestral allele. The dog may have a bit smaller size.

On behalf of Genoscoper Laboratories,

SIGNATURE



Nickname: Paavo Country: Finland

Registration ID: ER47004/19 Testing date: 2019/10/28

Microchip: 985141001209254

Breed: Cirneco dell'Etna

DNA Identified with standard identification ISAG 2006 markers profile:

Gender: Male

Dog's identity verified from microchip or tattoo by veterinarian or other authorised person during sample taking: Yes

Test results - Traits - page 4

Morphology

Trait	Genotype	Description
BMP3 c.1344C>A (p.Phe448Leu)	C/C	The dog does not carry the tested allele typically associated with shortened head (brachycephaly). The dog is more likely to have an elongated head (dolichocephaly).
SMOC2	D/D	The dog does not carry the tested allele typically associated with shortened head (brachycephaly). The dog is more likely to have an elongated head (dolichocephaly).
chr10:11072007	T/T	The dog does not carry an allele typically associated with floppy ears. The dog is more likely to have pricked than floppy ears.
T c.189C>G (p.lle63Met)	C/C	The dog does not carry the tested bobtail-causing genetic variant. The dog is most likely long-tailed.
EPAS1 (p.Gly305Ser)	G/G	The dog does not carry the tested variant associated with adaptation to high altitudes.
LIMBR1 DC-1	G/G	The dog does not carry the tested allele associated with hind dewclaws in Asian breeds. The dog is not likely to have hind dewclaws.
LIMBR1 DC-2	A/G	The dog carries one copy of the allele associated with hind dewclaws in western breeds. About 50% of the dogs with this genotype have hind dewclaws.
AXL4	D/D	The dog does not have the tested allele typically associated with blue eyes in Siberian Huskies. The dog is likely to have brown eyes.

On behalf of Genoscoper Laboratories,

SIGNATURE



Blood Disorders - page 1

Disorder	Mode of Inheritance	Result
Bleeding disorder due to P2RY12 defect	Autosomal Recessive	Clear
Canine Cyclic Neutropenia, Cyclic Hematopoiesis, Grey Collie Syndrome, (CN)	Autosomal Recessive	Clear
Canine Leukocyte Adhesion Deficiency (CLAD), type III	Autosomal Recessive	Clear
Canine Scott Syndrome, (CSS)	Autosomal Recessive	Clear
Factor IX Deficiency or Hemophilia B; mutation Gly379Glu	X-linked Recessive	Clear
Factor IX Deficiency or Hemophilia B; mutation originally found in Airedale Terrier	X-linked Recessive	Clear
Factor IX Deficiency or Hemophilia B; mutation originally found in Lhasa Apso	X-linked Recessive	Clear
Factor VII Deficiency	Autosomal Recessive	Clear
Factor VIII Deficiency or Hemophilia A; mutation originally found in Boxer	X-linked Recessive	Clear
Factor VIII Deficiency or Hemophilia A; mutation originally found in German Shepherd Dog	X-linked Recessive	Clear
Factor VIII Deficiency or Hemophilia A; mutation originally found in Old English Sheepdog	X-linked Recessive	Clear
Factor VIII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally found in German Shepherd	X-linked Recessive	Clear
Factor XI Deficiency	Autosomal Dominant (Incomplete Penetrance)	Clear
Familial Congenital Methemoglobinemia; mutation originally found in Pomeranian	Autosomal Recessive	Clear
Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Pyrenean Mountain Dog	Autosomal Recessive	Clear
Glanzmann Thrombasthenia Type I, (GT); mutation originally found in mixed breed dogs	Autosomal Recessive	Clear
Hereditary Elliptocytosis		Clear
Hereditary Phosphofructokinase (PFK) Deficiency	Autosomal Recessive	Clear
Macrothrombocytopenia; disease-linked variant originally found in Norfolk and Cairn Terrier	Autosomal Recessive	Clear
May-Hegglin Anomaly (MHA)	Autosomal Dominant	Clear
Prekallikrein Deficiency	Autosomal Recessive	Clear



Blood Disorders - page 2

Disorder	Mode of Inheritance	Result
Pyruvate Kinase Deficiency; mutation originally found in Basenji	Autosomal Recessive	Clear
Pyruvate Kinase Deficiency; mutation originally found in Beagle	Autosomal Recessive	Clear
Pyruvate Kinase Deficiency; mutation originally found in Pug	Autosomal Recessive	Clear
Pyruvate Kinase Deficiency; mutation originally found in West Highland White Terrier	Autosomal Recessive	Clear
Thrombopathia; mutation originally found in Basset Hound	Autosomal Recessive	Clear
Thrombopathia; mutation originally found in Eskimo Spitz	Autosomal Recessive	Clear
Thrombopathia; mutation originally found in Landseer	Autosomal Recessive	Clear
Trapped Neutrophil Syndrome, (TNS)	Autosomal Recessive	Clear
Von Willebrand's Disease (vWD) Type 1	Autosomal Recessive	Clear
Von Willebrand's Disease (vWD) Type 2	Autosomal Recessive	Clear
Von Willebrand's Disease (vWD) Type 3; mutation originally found in Kooikerhondje	Autosomal Recessive	Clear
Von Willebrand's Disease (vWD) Type 3; mutation originally found in Scottish Terrier	Autosomal Recessive	Clear
Von Willebrand's Disease (vWD) Type 3; mutation originally found in Shetland Sheepdog	Autosomal Recessive	Clear



Ocular Disorders - page 1

Disorder	Mode of Inheritance	Result
Canine Multifocal Retinopathy 1, (CMR1); mutation originally found in Mastiff-related breeds	Autosomal Recessive	Clear
Canine Multifocal Retinopathy 2, (CMR2); mutation originally found in Coton de Tulear	Autosomal Recessive	Clear
Canine Multifocal Retinopathy 3, (CMR3); mutation originally found in Lapponian Herder	Autosomal Recessive	Clear
Cone Degeneration, (CD) or Achromatopsia; mutation originally found in Alaskan Malamute	Autosomal Recessive	Clear
Cone Degeneration, (CD) or Achromatopsia; mutation originally found in German Shepherd Dog	Autosomal Recessive	Clear
Cone Degeneration, (CD) or Achromatopsia; mutation originally found in German Shorthaired Pointer	Autosomal Recessive	Clear
Cone-Rod Dystrophy 1, (crd1); mutation originally found in American Staffordshire Terrier	Autosomal Recessive	Clear
Cone-Rod Dystrophy 2, (crd2); mutation originally found in American Pit Bull Terrier	Autosomal Recessive	Clear
Cone-Rod Dystrophy, (cord1-PRA / crd4)	Autosomal Recessive (Incomplete Penetrance)	Clear
Cone-Rod Dystrophy, Standard Wirehaired Dachshund, (crd SWD)	Autosomal Recessive	Clear
Congenital Eye Disease; mutation originally found in Irish Soft-Coated Wheaten Terrier	Autosomal Recessive	Clear
Dominant Progressive Retinal Atrophy, (DPRA)	Autosomal Dominant	Clear
Early Retinal Degeneration, (erd); mutation originally found in Norwegian Elkhound	Autosomal Recessive	Clear
Generalized Progressive Retinal Atrophy	Autosomal Recessive	Clear
Golden Retriever Progressive Retinal Atrophy 1, (GR_PRA 1)	Autosomal Recessive	Clear
Golden Retriever Progressive Retinal Atrophy 2, (GR_PRA 2)	Autosomal Recessive	Clear
Primary Hereditary Cataract, (PHC); mutation originally found in Australian Shepherd	Autosomal Dominant (Incomplete Penetrance)	Clear
Primary Lens Luxation, (PLL)	Autosomal Recessive	Clear
Primary Open Angle Glaucoma, (POAG); mutation originally found in Basset Fauve de Bretagne	Autosomal Recessive	Clear
Primary Open Angle Glaucoma, (POAG); mutation originally found in Beagle	Autosomal Recessive	Clear
Primary Open Angle Glaucoma, (POAG); mutation originally found in Norwegian Elkhound	Autosomal Recessive	Clear



Ocular Disorders - page 2

Disorder	Mode of Inheritance	Result
Primary Open Angle Glaucoma, (POAG); mutation originally found in Petit Basset Griffon Vendeen	Autosomal Recessive	Clear
Primary lens luxation (PLL) and glaucoma; mutation originally found in Shar Pei	Autosomal Recessive	Clear
Progressive Retinal Atrophy (PRA4); mutation originally found in Lhasa Apso	Autosomal Recessive	Clear
Progressive Retinal Atrophy Type III, (PRA type III); mutation originally found in Tibetan Spaniel and Tibetan Terrier	Autosomal Recessive	Clear
Progressive Retinal Atrophy, (CNGA1-PRA); mutation originally found in Shetland Sheepdog	Autosomal Recessive	Clear
Progressive Retinal Atrophy, (PAP1_PRA); mutation originally found in Papillon and Phalene	Autosomal Recessive	Clear
Progressive Retinal Atrophy, (PRA); mutation originally found in Basenji	Autosomal Recessive	Clear
Progressive Retinal Atrophy, (PRA); mutation originally found in Swedish Vallhund	Autosomal Recessive	Clear
Rod-Cone Dysplasia 1, (rcd1); mutation originally found in Irish Setter	Autosomal Recessive	Clear
Rod-Cone Dysplasia 1a, (rdc1a); mutation originally found in Sloughi	Autosomal Recessive	Clear
Rod-Cone Dysplasia 3, (rcd3)	Autosomal Recessive	Clear
X-Linked Progressive Retinal Atrophy 1, (XLPRA1)	X-linked Recessive	Clear
X-Linked Progressive Retinal Atrophy 2, (XLPRA2; Type A PRA)	X-linked Recessive	Clear

Cardiac Disorders

Disorder	Mode of Inheritance	Result
Dilated Cardiomyopathy, (DCM); mutation originally found in Schnauzer	Autosomal Recessive	Clear
Long QT Syndrome	Autosomal Dominant	Clear



Endocrine Disorders

Disorder	Mode of Inheritance	Result
Congenital Hypothyroidism; mutation originally found in Tenterfield Terrier	Autosomal Recessive	Clear
Congenital Hypothyroidism; mutation originally found in Toy Fox and Rat Terrier	Autosomal Recessive	Clear

Immunological Disorders

Disorder	Mode of Inheritance	Result
Autosomal Recessive Severe Combined Immunodeficiency, (ARSCID)	Autosomal Recessive	Clear
Complement 3 (C3) Deficiency	Autosomal Recessive	Clear
Myeloperoxidase Deficiency	Autosomal Recessive	Clear
Severe Combined Immunodeficiency in Frisian Water Dogs, (SCID)	Autosomal Recessive	Clear
X-Linked Severe Combined Immunodeficiency (XSCID); mutation originally found in Basset Hound	X-linked Recessive	Clear
X-Linked Severe Combined Immunodeficiency (XSCID); mutation originally found in Cardigan Welsh Corgi	X-linked Recessive	Clear



Renal Disorders

Disorder	Mode of Inheritance	Result
2,8-Dihydroxyadenine (2,8-DHA) urolithiasis	Autosomal Recessive	Clear
Cystinuria Type I-A; mutation originally found in Newfoundland Dog	Autosomal Recessive	Clear
Cystinuria Type II-A; mutation originally found in Australian Cattle Dog	Autosomal Dominant	Clear
Familial Nephropathy (FN); mutation originally found in English Cocker Spaniel	Autosomal Recessive	Clear
Familial Nephropathy (FN); mutation originally found in English Springer Spaniel	Autosomal Recessive	Clear
Fanconi Syndrome	Autosomal Recessive	Clear
Hyperuricosuria, (HUU)	Autosomal Recessive	Clear
Polycystic Kidney Disease in Bull Terriers, (BTPKD)	Autosomal Dominant	Clear
Primary Hyperoxaluria, (PH); mutation originally found in Coton de Tulear	Autosomal Recessive	Clear
Protein Losing Nephropathy, (PLN); NPHS1 gene variant		Clear
Renal Cystadenocarcinoma and Nodular Dermatofibrosis, (RCND)	Autosomal Dominant	Clear
X-Linked Hereditary Nephropathy, (XLHN)	X-linked Recessive	Clear
X-Linked Hereditary Nephropathy, (XLHN); mutation originally found in Navasota Dog	X-linked Recessive	Clear
Xanthinuria, Type 1a; mutation originally found in mixed breed dogs	Autosomal Recessive	Clear
Xanthinuria, Type 2a; mutation originally found in Toy Manchester Terrier	Autosomal Recessive	Clear
Xanthinuria, Type 2b; mutation originally found in Cavalier King Charles Spaniel and English Cocker Spaniel	Autosomal Recessive	Clear



Metabolic Disorders

Disorder	Mode of Inheritance	Result
Glycogen Storage Disease Type II or Pompe's Disease, (GSD II)	Autosomal Recessive	Clear
Glycogen Storage Disease Type IIIa, (GSD IIIa)	Autosomal Recessive	Clear
Glycogen Storage Disease Type Ia, (GSD Ia)	Autosomal Recessive	Clear
Hypocatalasia or Acatalasemia	Autosomal Recessive	Clear
Intestinal Cobalamin Malabsorption or Imerslund-Gräsbeck Syndrome, (IGS); mutation originally found in Beagle	Autosomal Recessive	Clear
Intestinal Cobalamin Malabsorption or Imerslund-Gräsbeck Syndrome, (IGS); mutation originally found in Border Collie	Autosomal Recessive	Clear
Mucopolysaccharidosis Type IIIA, (MPS IIIA); mutation originally found in Dachshund	Autosomal Recessive	Clear
Mucopolysaccharidosis Type IIIA, (MPS IIIA); mutation originally found in New Zealand Huntaway	Autosomal Recessive	Clear
Mucopolysaccharidosis Type VII, (MPS VII); mutation originally found in Brazilian Terrier	Autosomal Recessive	Clear
Mucopolysaccharidosis Type VII, (MPS VII); mutation originally found in German Shepherd	Autosomal Recessive	Clear
Pyruvate Dehydrogenase Phosphatase 1 (PDP1) Deficiency	Autosomal Recessive	Clear



Muscular Disorders

Disorder	Mode of Inheritance	Result
Cavalier King Charles Spaniel Muscular Dystrophy, (CKCS-MD)	X-linked Recessive	Clear
Centronuclear Myopathy, (CNM); mutation originally found in Great Dane	Autosomal Recessive	Clear
Centronuclear Myopathy, (CNM); mutation originally found in Labrador Retriever	Autosomal Recessive	Clear
Duchenne or Dystrophin Muscular Dystrophy, (DMD); mutation originally found in Golden Retriever	X-linked Recessive	Clear
Duchenne or Dystrophin Muscular Dystrophy, (DMD); mutation originally found in Norfolk Terrier	X-linked Recessive	Clear
Muscular Dystrophy, Ullrich-type; mutation originally found in Landseer	Autosomal Recessive	Clear
Myostatin deficiency (Double Muscling, "Bully")	Autosomal Recessive	Clear
Myotonia Congenita; mutation originally found in Australian Cattle Dog	Autosomal Recessive	Clear
Myotonia Congenita; mutation originally found in in Labrador Retriever	Autosomal Recessive	Clear
Myotubular Myopathy; mutation originally found in Rottweiler	X-linked Recessive	Clear
Nemaline Myopathy; mutation originally found in American Bulldog	Autosomal Recessive	Clear
X-Linked Myotubular Myopathy	X-linked Recessive	Clear



Neurological Disorders - page 1

Alaskan Husky Encephalopathy, (AHE) Autosomal Recessive Clear Alexander Disease (AxD); mutation originally found in Labrador Retriever Autosomal Dominant Clear Bandera's Neonatal Ataxia, (BNAt) Benign Familial Juvenile Epilepsy or Remitting Focal Epilepsy Cerebellar Cortical Degeneration, (CCD); mutation originally found in Vizsla Cerebral Dysfunction; mutation originally found in Friesian Stabyhoun Autosomal Recessive Clear Clear Dandy-Walker-Like Malformation (DWLM); mutation originally found in Eurasier Degenerative Myelopathy, (DM; SOD1A) Autosomal Recessive (Incomplete Penetrance) Clear Clear Autosomal Recessive Clear Autosomal Recessive Clear Clear Autosomal Recessive Clear Clear Autosomal Recessive Clear Clear Clear Autosomal Recessive Clear Clear Autosomal Recessive Clear Clear Clear Autosomal Recessive Clear C	Disorder	Mode of Inheritance	Result
Altosomal Dominant Clear Bandera's Neonatal Ataxia, (BNAt) Benign Familial Juvenile Epilepsy or Remitting Focal Epilepsy Benign Familial Juvenile Epilepsy Benign Familial Juvenile Polymeurolation originally found in Benign Familial Juvenile Autosomal Recessive Clear Benign Familial Juvenile Autosomal Recessive Clear Benign Familial Juvenile Myoclonic Epilepsy, (JME); mutation originally found in Benign Familial Juvenile Myoclonic Epilepsy, (JME); mutation originally found in Benign Familial Juvenile Autosomal Recessive Clear Benign Familial Juvenile Myoclonic Epilepsy, (JME); mutation originally found in Benign Familial Juvenile Myoclonic Epilepsy, (JME); mutation originally found in Benign Familial Juvenile Benessive Clear Benign Familial Familia Parson Russell Clear Benign Familial Pacessive Clear Benign Fami	Acral Mutilation Syndrome, (AMS)	Autosomal Recessive	Clear
Bandera's Neonatal Ataxia, (BNAt) Benign Familial Juvenile Epilepsy or Remitting Focal Epilepsy Benign Familial Juvenile Epilepsy or Remitting Focal Epilepsy Autosomal Recessive Clear Cerebellar Cortical Degeneration, (CCD); mutation originally found in Vizsla Cerebral Dysfunction; mutation originally found in Friesian Stabyhoun Autosomal Recessive Clear Dendy-Walker-Like Malformation (DWLM); mutation originally found in Eurasier Degenerative Myelopathy, (DM; SOD1A) Clear Early-Onset Progressive Polyneuropathy; mutation originally found in Autosomal Recessive (Incomplete Penetrance) Early-Onset Progressive Polyneuropathy; mutation originally found in Autosomal Recessive Clear Early-Onset Progressive Polyneuropathy; mutation originally found in Autosomal Recessive Clear Hereditary Ataxia or Cerebellar Ataxia; mutation originally found in Old English Sheepdog and Gordon Setter Hereditary Ataxia; mutation originally found in in Norwegian Buhund Autosomal Recessive Clear Hyperekplexia or Startle Disease Autosomal Recessive Clear Hyperekplexia or Startle Disease Autosomal Recessive Clear Hypomyelination; mutation originally found in Weimaraner Autosomal Recessive Clear Hypomyelination; mutation originally found in Weimaraner Autosomal Recessive Clear Hypomyelination; mutation originally found in Parson Russell Autosomal Recessive Clear Clear L'2-Hydroxyglutaric aciduria, (L2HGA); mutation originally found in Battordshire Bull Terrier L-2-Hydroxyglutaric aciduria, (L2HGA); mutation originally found in West Highland White Terrier	Alaskan Husky Encephalopathy, (AHE)	Autosomal Recessive	Clear
Benign Familial Juvenile Epilepsy or Remitting Focal Epilepsy Autosomal Recessive Clear Cerebellar Cortical Degeneration, (CCD); mutation originally found in Autosomal Recessive Clear Cerebral Dysfunction; mutation originally found in Friesian Stabyhoun Autosomal Recessive Clear Dandy-Walker-Like Malformation (DWLM); mutation originally found in Eurasier Degenerative Myelopathy, (DM; SOD1A) Autosomal Recessive (Incomplete Penetrance) Early-Onset Progressive Polyneuropathy; mutation originally found in Autosomal Recessive Clear Autosomal Recessive Clear Autosomal Recessive Clear Autosomal Recessive Clear Fetal Onset Progressive Polyneuropathy; mutation originally found in Greyhound Fetal Onset Neuroaxonal Dystrophy, (FNAD) Autosomal Recessive Clear Hereditary Ataxia or Cerebellar Ataxia; mutation originally found in Old English Sheepdog and Gordon Setter Hereditary Ataxia; mutation originally found in in Norwegian Buhund Autosomal Recessive Clear Hypomyelination; mutation originally found in Weimaraner Autosomal Recessive Clear Juvenile Myoclonic Epilepsy, (JME); mutation originally found in Autosomal Recessive Clear Autosomal Recessive Clear Juvenile Myoclonic Epilepsy, (JME); mutation originally found in Parson Russell Autosomal Recessive Clear Autosomal Recessive Clear L-2-Hydroxyglutaric aciduria, (L2HGA); mutation originally found in West Highland White Terrier Autosomal Recessive Clear	Alexander Disease (AxD); mutation originally found in Labrador Retriever	Autosomal Dominant	Clear
Cerebellar Cortical Degeneration, (CCD); mutation originally found in Vizsla Cerebral Dysfunction; mutation originally found in Friesian Stabyhoun Autosomal Recessive Clear Dandy-Walker-Like Malformation (DWLM); mutation originally found in Autosomal Recessive Clear Degenerative Myelopathy, (DM; SOD1A) Early-Onset Progressive Polyneuropathy; mutation originally found in Autosomal Recessive (Incomplete Penetrance) Early-Onset Progressive Polyneuropathy; mutation originally found in Autosomal Recessive Clear Greyhound Early-Onset Progressive Polyneuropathy; mutation originally found in Autosomal Recessive Clear Greyhound Fetal Onset Neuroaxonal Dystrophy, (FNAD) Hereditary Ataxia or Cerebellar Ataxia; mutation originally found in Old English Sheepdog and Gordon Setter Hereditary Ataxia; mutation originally found in Norwegian Buhund Autosomal Recessive Clear Hyporekplexia or Startle Disease Autosomal Recessive Clear Autosomal Recessive Clear Hypomyelination; mutation originally found in Weimaraner Autosomal Recessive Clear Duvenile Myoclonic Epilepsy, (JME); mutation originally found in Autosomal Recessive Clear Clear Phypomyelination; mutation originally found in Parson Russell Autosomal Recessive Clear Clear Duvenile encephalopathy; mutation originally found in Parson Russell Autosomal Recessive Clear Clear Duvenile encephalopathy; mutation originally found in Parson Russell Autosomal Recessive Clear Clear Duvenile encephalopathy; mutation originally found in Parson Russell Clear Clear Duvenile encephalopathy; mutation originally found in Parson Russell Clear C	Bandera's Neonatal Ataxia, (BNAt)	Autosomal Recessive	Clear
Cerebral Dysfunction; mutation originally found in Friesian Stabyhoun Autosomal Recessive Clear Dandy-Walker-Like Malformation (DWLM); mutation originally found in Eurasier Degenerative Myelopathy, (DM; SOD1A) Autosomal Recessive Clear Autosomal Recessive (Incomplete Penetrance) Clear Autosomal Recessive Clear Clear Autosomal Recessive Clear Autosomal Recessive Clear Hereditary Ataxia or Cerebellar Ataxia; mutation originally found in Old English Sheepdog and Gordon Setter Hereditary Ataxia; mutation originally found in Norwegian Buhund Autosomal Recessive Clear Hyperekplexia or Startle Disease Autosomal Recessive Clear Hypomyelination; mutation originally found in Weimaraner Autosomal Recessive Clear Autosomal Recessive Clear Juvenile Myoclonic Epilepsy, (JME); mutation originally found in Rhodesian Ridgeback Lear Lear-Hydroxyglutaric aciduria, (L2HGA); mutation originally found in Staffordshire Bull Terrier Lear-Hydroxyglutaric aciduria, (L2HGA); mutation originally found in West Highland White Terrier	Benign Familial Juvenile Epilepsy or Remitting Focal Epilepsy	Autosomal Recessive	Clear
Dandy-Walker-Like Malformation (DWLM); mutation originally found in Degenerative Myelopathy, (DM; SOD1A) Autosomal Recessive (Incomplete Penetrance) Early-Onset Progressive Polyneuropathy; mutation originally found in Alaskan Malamute Early-Onset Progressive Polyneuropathy; mutation originally found in Autosomal Recessive Clear Greyhound Autosomal Recessive Clear Clear Greyhound Clear Clear Clear Autosomal Recessive Clear Autosomal Recessive Clear Clear Clear Autosomal Recessive Clear Autosomal Recessive Clear C	Cerebellar Cortical Degeneration, (CCD); mutation originally found in Vizsla	Autosomal Recessive	Clear
Eurasier Degenerative Myelopathy, (DM; SOD1A) Autosomal Recessive (Incomplete Penetrance) Early-Onset Progressive Polyneuropathy; mutation originally found in Autosomal Recessive Clear Alaskan Malamute Early-Onset Progressive Polyneuropathy; mutation originally found in Autosomal Recessive Clear Greyhound Fetal Onset Neuroaxonal Dystrophy, (FNAD) Hereditary Ataxia or Cerebellar Ataxia; mutation originally found in Old Hereditary Ataxia or Cerebellar Ataxia; mutation originally found in Old Hereditary Ataxia; mutation originally found in Norwegian Buhund Autosomal Recessive Clear Hyperekplexia or Startle Disease Hyperekplexia or Startle Disease Hypomyelination; mutation originally found in Weimaraner Autosomal Recessive Clear Hypomyelination; mutation originally found in Autosomal Recessive Clear Hypomyelination; mutation originally found in Parson Russell Duvenile Myoclonic Epilepsy, (JME); mutation originally found in Autosomal Recessive Clear Clear Clear Autosomal Recessive Clear Clear Clear Autosomal Recessive Clear	Cerebral Dysfunction; mutation originally found in Friesian Stabyhoun	Autosomal Recessive	Clear
Early-Onset Progressive Polyneuropathy; mutation originally found in Autosomal Recessive Clear Alaskan Malamute Early-Onset Progressive Polyneuropathy; mutation originally found in Greyhound Autosomal Recessive Clear Greyhound Fetal Onset Neuroaxonal Dystrophy, (FNAD) Autosomal Recessive Clear Hereditary Ataxia or Cerebellar Ataxia; mutation originally found in Old Autosomal Recessive Clear English Sheepdog and Gordon Setter Hereditary Ataxia; mutation originally found in Norwegian Buhund Autosomal Recessive Clear Hyperekplexia or Startle Disease Autosomal Recessive Clear Hypomyelination; mutation originally found in Weimaraner Autosomal Recessive Clear Duvenile Myoclonic Epilepsy, (JME); mutation originally found in Autosomal Recessive Clear Phyonesian Ridgeback Juvenile encephalopathy; mutation originally found in Parson Russell Autosomal Recessive Clear Terrier L-2-Hydroxyglutaric aciduria, (L2HGA); mutation originally found in West Autosomal Recessive Clear Clear Staffordshire Bull Terrier L-2-Hydroxyglutaric aciduria, (L2HGA); mutation originally found in West Autosomal Recessive Clear Clear Staffordshire Bull Terrier Clear Autosomal Recessive Clear Cl	Dandy-Walker-Like Malformation (DWLM); mutation originally found in Eurasier	Autosomal Recessive	Clear
Alaskan Malamute Early-Onset Progressive Polyneuropathy; mutation originally found in Greyhound Fetal Onset Neuroaxonal Dystrophy, (FNAD) Hereditary Ataxia or Cerebellar Ataxia; mutation originally found in Old Autosomal Recessive Hereditary Ataxia; mutation originally found in Norwegian Buhund Hereditary Ataxia; mutation originally found in in Norwegian Buhund Autosomal Recessive Clear Hyperekplexia or Startle Disease Autosomal Recessive Clear Hypomyelination; mutation originally found in Weimaraner Autosomal Recessive Clear Juvenile Myoclonic Epilepsy, (JME); mutation originally found in Rhodesian Ridgeback Juvenile encephalopathy; mutation originally found in Parson Russell Terrier L-2-Hydroxyglutaric aciduria, (L2HGA); mutation originally found in Staffordshire Bull Terrier Clear Clear Clear Autosomal Recessive Clear	Degenerative Myelopathy, (DM; SOD1A)		Clear
Fetal Onset Neuroaxonal Dystrophy, (FNAD) Autosomal Recessive Clear Hereditary Ataxia or Cerebellar Ataxia; mutation originally found in Old English Sheepdog and Gordon Setter Hereditary Ataxia; mutation originally found in in Norwegian Buhund Autosomal Recessive Clear Hyperekplexia or Startle Disease Autosomal Recessive Clear Hypomyelination; mutation originally found in Weimaraner Autosomal Recessive Clear Juvenile Myoclonic Epilepsy, (JME); mutation originally found in Rhodesian Ridgeback Juvenile encephalopathy; mutation originally found in Parson Russell Terrier L-2-Hydroxyglutaric aciduria, (L2HGA); mutation originally found in Staffordshire Bull Terrier Clear Clear Autosomal Recessive Clear Clear Autosomal Recessive Clear Autosomal Recessive Clear Autosomal Recessive Clear Clear Autosomal Recessive Clear Clear Autosomal Recessive Clear Clear Autosomal Recessive Clear Autosomal Recessive Clear Autosomal Recessive Clear Clear Autosomal Recessive Clear	Early-Onset Progressive Polyneuropathy; mutation originally found in Alaskan Malamute	Autosomal Recessive	Clear
Hereditary Ataxia or Cerebellar Ataxia; mutation originally found in Old English Sheepdog and Gordon Setter Hereditary Ataxia; mutation originally found in in Norwegian Buhund Autosomal Recessive Clear Hyperekplexia or Startle Disease Autosomal Recessive Clear Hypomyelination; mutation originally found in Weimaraner Autosomal Recessive Clear Hypomyelination; mutation originally found in Weimaraner Autosomal Recessive Clear Clear Hupomyelination; mutation originally found in Parson Russell Clear Rhodesian Ridgeback Juvenile encephalopathy; mutation originally found in Parson Russell Clear Terrier L-2-Hydroxyglutaric aciduria, (L2HGA); mutation originally found in Staffordshire Bull Terrier Clear Autosomal Recessive Clear Clear Autosomal Recessive Clear Clear Autosomal Recessive Clear Clear Autosomal Recessive Clear	Early-Onset Progressive Polyneuropathy; mutation originally found in Greyhound	Autosomal Recessive	Clear
English Sheepdog and Gordon Setter Hereditary Ataxia; mutation originally found in in Norwegian Buhund Autosomal Recessive Clear Hyperekplexia or Startle Disease Autosomal Recessive Clear Hypomyelination; mutation originally found in Weimaraner Autosomal Recessive Clear Juvenile Myoclonic Epilepsy, (JME); mutation originally found in Rhodesian Ridgeback Juvenile encephalopathy; mutation originally found in Parson Russell Autosomal Recessive Clear Terrier L-2-Hydroxyglutaric aciduria, (L2HGA); mutation originally found in Autosomal Recessive Clear Staffordshire Bull Terrier L-2-Hydroxyglutaric aciduria, (L2HGA); mutation originally found in West Highland White Terrier Clear	Fetal Onset Neuroaxonal Dystrophy, (FNAD)	Autosomal Recessive	Clear
Hyperekplexia or Startle Disease Autosomal Recessive Clear Hypomyelination; mutation originally found in Weimaraner Autosomal Recessive Clear Juvenile Myoclonic Epilepsy, (JME); mutation originally found in Rhodesian Ridgeback Juvenile encephalopathy; mutation originally found in Parson Russell Terrier L-2-Hydroxyglutaric aciduria, (L2HGA); mutation originally found in Staffordshire Bull Terrier L-2-Hydroxyglutaric aciduria, (L2HGA); mutation originally found in West Highland White Terrier Clear	Hereditary Ataxia or Cerebellar Ataxia; mutation originally found in Old English Sheepdog and Gordon Setter	Autosomal Recessive	Clear
Hypomyelination; mutation originally found in Weimaraner Autosomal Recessive Clear Glavenile Myoclonic Epilepsy, (JME); mutation originally found in Rhodesian Ridgeback Juvenile encephalopathy; mutation originally found in Parson Russell Autosomal Recessive Clear	Hereditary Ataxia; mutation originally found in in Norwegian Buhund	Autosomal Recessive	Clear
Juvenile Myoclonic Epilepsy, (JME); mutation originally found in Rhodesian Ridgeback Juvenile encephalopathy; mutation originally found in Parson Russell Terrier L-2-Hydroxyglutaric aciduria, (L2HGA); mutation originally found in Staffordshire Bull Terrier L-2-Hydroxyglutaric aciduria, (L2HGA); mutation originally found in West Highland White Terrier Clear	Hyperekplexia or Startle Disease	Autosomal Recessive	Clear
Rhodesian Ridgeback Juvenile encephalopathy; mutation originally found in Parson Russell L-2-Hydroxyglutaric aciduria, (L2HGA); mutation originally found in Staffordshire Bull Terrier L-2-Hydroxyglutaric aciduria, (L2HGA); mutation originally found in West Highland White Terrier Autosomal Recessive Clear Clear	Hypomyelination; mutation originally found in Weimaraner	Autosomal Recessive	Clear
Terrier L-2-Hydroxyglutaric aciduria, (L2HGA); mutation originally found in Staffordshire Bull Terrier L-2-Hydroxyglutaric aciduria, (L2HGA); mutation originally found in West Highland White Terrier Autosomal Recessive Clear	Juvenile Myoclonic Epilepsy, (JME); mutation originally found in Rhodesian Ridgeback	Autosomal Recessive	Clear
Staffordshire Bull Terrier L-2-Hydroxyglutaric aciduria, (L2HGA); mutation originally found in West Autosomal Recessive Clear Highland White Terrier	Juvenile encephalopathy; mutation originally found in Parson Russell Terrier	Autosomal Recessive	Clear
Highland White Terrier	L-2-Hydroxyglutaric aciduria, (L2HGA); mutation originally found in Staffordshire Bull Terrier	Autosomal Recessive	Clear
_agotto Storage Disease, (LSD) Autosomal Recessive Clear	L-2-Hydroxyglutaric aciduria, (L2HGA); mutation originally found in West Highland White Terrier	Autosomal Recessive	Clear
	Lagotto Storage Disease, (LSD)	Autosomal Recessive	Clear



Neurological Disorders - page 2

Disorder	Mode of Inheritance	Result	
Neonatal Cerebellar Cortical Degeneration or Cerebellar Abiotrophy, (NCCD)	Autosomal Recessive	Clear	
Neonatal Encephalopathy with Seizures, (NEWS)	Autosomal Recessive	Clear	
Neuroaxonal Dystrophy (NAD); mutation originally found in Spanish Water Dog	Autosomal Recessive	Clear	
Neuroaxonal Dystrophy, (NAD); mutation originally found in Papillon	Autosomal Recessive	Clear	
Neuronal Ceroid Lipofuscinosis 1, (NCL1); mutation originally found in Dachshund	Autosomal Recessive	Clear	
Neuronal Ceroid Lipofuscinosis 10, (NCL10); mutation originally found in American Bulldog	Autosomal Recessive	Clear	
Neuronal Ceroid Lipofuscinosis 4A, (NCL4); mutation originally found in American Staffordshire Terrier	Autosomal Recessive	Clear	
Neuronal Ceroid Lipofuscinosis 5, (NCL5); mutation originally found in Border Collie	Autosomal Recessive	Clear	
Neuronal Ceroid Lipofuscinosis 8, (NCL8); mutation originally found in Alpine Dachsbracke	Autosomal Recessive	Clear	
Neuronal Ceroid Lipofuscinosis 8, (NCL8); mutation originally found in Australian Shepherd	Autosomal Recessive	Clear	
Neuronal Ceroid Lipofuscinosis 8, (NCL8); mutation originally found in English Setter	Autosomal Recessive	Clear	
Neuronal Ceroid Lipofuscinosis, (NCL7); mutation originally found in Chinese Crested Dog and Chihuahua	Autosomal Recessive	Clear	
Polyneuropathy with ocular abnormalities and neuronal vacuolation, (POANV); mutation originally found in Black Russian Terrier	Autosomal Recessive	Clear	
Progressive Early-Onset Cerebellar Ataxia; mutation originally found in Finnish Hound	Autosomal Recessive	Clear	
Sensory Neuropathy; mutation originally found in Border Collie	Autosomal Recessive	Clear	
Shaking Puppy Spongiform LeucoEncephaloMyelopathy, (SLEM); mutation originally found in Border Terrier	Autosomal Recessive	Clear	
Spinocerebellar Ataxia with Myokymia and/or Seizures (SCA)	Autosomal Recessive	Clear	
Spinocerebellar Ataxia/ Late-Onset Ataxia (SCA, LOA)	Autosomal Recessive	Clear	
Spongy Degeneration with Cerebellar Ataxia, (SDCA1); mutation originally found in Belgian Shepherd Dog	Autosomal Recessive	Clear	
Spongy Degeneration with Cerebellar Ataxia, (SDCA2); mutation originally found in Belgian Shepherd Dog	Autosomal Recessive	Clear	
X-Linked Tremors; mutation originally found in English Springer Spaniel	X-linked Recessive	Clear	



Neuromuscular Disorders

Disorder	Mode of Inheritance	Result	
Congenital Myasthenic Syndrome (CMS); mutation originally found in Labrador Retriever	Autosomal Recessive	Clear	
Congenital Myasthenic Syndrome, (CMS); mutation originally found in Jack Russell Terrier	Autosomal Recessive	Clear	
Congenital Myasthenic Syndrome, (CMS); mutation originally found in Old Danish Pointing Dog	Autosomal Recessive Clea		
Exercise-Induced Collapse, (EIC)	Autosomal Recessive (Incomplete Penetrance)	Clear	
GM1 Gangliosidosis; mutation originally found in Alaskan Husky	Autosomal Recessive	Clear	
GM1 Gangliosidosis; mutation originally found in Portuguese Water Dog	Autosomal Recessive	Clear	
GM1 Gangliosidosis; mutation originally found in Shiba Dog	Autosomal Recessive	Clear	
GM2 Gangliosidosis, mutation originally found in Japanese Chin	Autosomal Recessive	Clear	
GM2 Gangliosidosis; mutation originally found in Toy Poodle	Autosomal Recessive	Clear	
Globoid Cell Leukodystrophy or Krabbe Disease, (GLD); mutation originally found in Irish Setter	Autosomal Recessive	Clear	
Globoid Cell Leukodystrophy or Krabbe Disease, (GLD); mutation originally found in Terriers	Autosomal Recessive	Clear	
Paroxysmal Dyskinesia, (PxD); mutation originally found in Irish Soft Coated Wheaten Terrier	Autosomal Recessive	Clear	



Skeletal Disorders

Chondrodysplasia; mutation originally found in Norwegian Elkhound and Karelian Bear Dog	Autosomal Recessive	Clear			
Cleft Palate; Cleft Lip and Palate with Syndactyly; ADAMTS20 gene mutation originally found in Nova Scotia Duck Tolling Retriever	Autosomal Recessive	Clear			
Cleft Palate; DLX6 gene mutation originally found in Nova Scotia Duck Tolling Retriever	Autosomal Recessive	Clear			
Craniomandibular Osteopathy, (CMO); mutation associated with terrier breeds	Autosomal Dominant Clear (Incomplete Penetrance)				
Hereditary Vitamin D-Resistant Rickets, (HVDRR)	Autosomal Recessive	Clear			
Oculoskeletal Dysplasia 2 or Dwarfism-Retinal Dysplasia 2, (OSD2)	Autosomal Recessive	Clear			
Osteochondrodysplasia; mutation originally found in Miniature Poodle	Autosomal Recessive	Clear			
Osteochondromatosis; mutation originally found in American Staffordshire Terrier	Autosomal Dominant	Clear			
Osteogenesis Imperfecta, (OI); mutation originally found in Beagle	Autosomal Dominant	Clear			
Osteogenesis Imperfecta, (OI); mutation originally found in Dachshund	Autosomal Recessive	Clear			
Skeletal Disease (Hypophosphatasia); mutation originally found in Karelian Bear Dog	Autosomal Recessive	Clear			
Skeletal Dysplasia 2, (SD2)	Autosomal Recessive	Clear			
Spondylocostal Dysostosis	Autosomal Recessive	Clear			
Van den Ende-Gupta Syndrome, (VDEGS)	Autosomal Recessive	Clear			



Dermal Disorders

Disorder	Mode of Inheritance	Result			
Dystrophic Epidermolysis Bullosa; mutation originally found in Central Asian Ovcharka	Autosomal Recessive	Clear			
Dystrophic Epidermolysis Bullosa; mutation originally found in Golden Retriever	Autosomal Recessive	Clear			
Epidermolytic Hyperkeratosis	Autosomal Recessive	Clear			
Focal Non-Epidermolytic Palmoplantar Keratoderma, (FNEPPK); mutation originally found in Dogue de Bordeaux	Autosomal Recessive	Clear			
Golden Retriever Ichthyosis	Autosomal Recessive	Clear			
Hereditary Footpad Hyperkeratosis, (HFH)	Autosomal Recessive	Clear			
Hereditary Nasal Parakeratosis, (HNPK); mutation originally found in Greyhound	Autosomal Recessive	Clear			
Ichthyosis; mutation originally found in American Bulldog	Autosomal Recessive	Clear			
Ichthyosis; mutation originally found in Great Dane	Autosomal Recessive	Clear			
Lamellar Ichthyosis, (LI)	Autosomal Recessive	Clear			
Lethal Acrodermatitis, (LAD); mutation originally found in in Bull Terrier and Miniature Bull Terrier	Autosomal Recessive	Clear			
Ligneous Membranitis	Autosomal Recessive	Clear			
Musladin-Lueke syndrome, (MLS)	Autosomal Recessive	Clear			
X-Linked Ectodermal Dysplasia, (XHED)	X-linked Recessive	Clear			



Other Disorders

Disorder	Mode of Inheritance	Result
Acute Respiratory Distress Syndrome, (ARDS); mutation originally found in Dalmatian	Autosomal Recessive	Clear
Amelogenesis Imperfecta, (AI); mutation originally found in Italian Greyhound	Autosomal Recessive	Clear
Amelogenesis Imperfecta, (AI); mutation originally found in Parson Russell Terrier	Autosomal Recessive	Clear
Congenital Keratoconjunctivitis Sicca and Ichthyosiform Dermatosis, (CKCSID)	Autosomal Recessive	Clear
Dental Hypomineralisation; mutation originally found in Border Collie	Autosomal Recessive	Clear
Lung Developmental Disease; mutation originally found in in Airedale Terrier	Autosomal Recessive	Clear
Narcolepsy; mutation originally found in Dachshund	Autosomal Recessive	Clear
Narcolepsy; mutation originally found in Doberman Pinscher	Autosomal Recessive	Clear
Narcolepsy; mutation originally found in Labrador Retriever	Autosomal Recessive	Clear
Persistent Müllerian Duct Syndrome, (PMDS); mutation originally found in Miniature Schnauzer	Autosomal Recessive	Clear
Primary Ciliary Dyskinesia, (PCD)	Autosomal Recessive	Clear



APPENDIX Explanation of the results of the tested disorders

Autosomal recessive inheritance (ARI)

Clear - A dog carries no copies of the tested mutation and has no or reduced likelihood of developing and passing on the disease/condition.

Carrier - A dog carries one copy of the tested mutation. Carriers typically have a normal, healthy appearance but pass on the mutation to approximately 50% of their offspring.

At risk - A dog carries two copies of the tested mutation and is at high or increased risk of developing the disease/condition.

Autosomal dominant inheritance (ADI)

Clear - A dog carries no copies of the tested mutation and has no or reduced likelihood of developing and passing on the disease/condition.

At risk - A dog carries one or two copies of the tested mutation and is at high or increased risk of developing the disease/condition.

X-linked recessive inheritance (X-linked)

Clear - A dog carries no copies of the tested mutation and has no or reduced likelihood of developing and passing on the disease/condition.

Carrier - Female carriers typically have a normal, healthy appearance but carry one copy of the tested mutation on one of their X chromosomes. As males only have one X chromosome, there are no male carriers.

At risk - Female dogs at risk carry two mutated copies of the tested mutation. Males carry one copy of the tested mutation on their single X chromosome. Dogs at risk are at high or increased risk of developing the disease/condition.

Please note that the descriptions above are generalized based on typically observed inheritance patterns. When obtaining a 'carrier' or 'at risk' test result, always refer to the corresponding online test documentation for more detailed information on the condition and any exceptions.

Genoscoper Laboratories - Legal Notice

Genoscoper Laboratories' services and test results are produced based on samples and materials supplied by the Client. Testing and analysis is performed by using methods and processes that Genoscoper Laboratories deems appropriate. Genoscoper Laboratories reserves the right to make changes in the collection of the single-gene tests included in the testing service as well as to remove results derived from them, if new information comes available that in any way questions the validity of the test results. Results provided by Genoscoper Laboratories are prepared solely for the use of the Client. For further information, please visit: www.mydogdna.com/legal-notices

Registered Name: Trucker's Fuel To The Fire

Nickname: Pauvo Registration ID: ER47004/19 Microchip: 985141001209254 Breed: Cirneco dell'Etna Gender: Male

Owner: Sanna Koponen
Country: Finland
Testing date: 28/10/2019
DNA identification profile:
Identified with standard ISAG 2006 markers
Dog's identity verified from microchip or tattoo by veterinarian or other authorised person during sample taking: Yes

PROVIDED BY GENOSCOPER LABORATORIES





Certificate of DNA Identification Profile

15/8/2020

REN64E19 149/153	REN247M23 266/268	REN169O18 170/170	REN169D01 216/216	 INU030 146/150	126/126	FH2848 230/240	AMELOGENIN Y/X	AHT137 147/147	REN54P11 228/232
REN162C04 204/204	REN105LO3 237/241	INRA21 95/103	FH2054 148/168	 AHTK253 288/288		AHTH171 221/233	AHTH130 119/119	AHTK211 87/91	AHT121 96/102

On behalf of Genoscoper Laboratories,

Jonas Donner, PhD, Head of Research and Development at Genoscoper Laboratories

Joras Domer

PO Box. 1040 FI-00251 Helsinki | info@genoscoper.com